

PCTWORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification: C12N 15/82, A01H 5/00, A01H 5/06	A1	(11) International Publication Number: WO 00/44919 (43) International Publication Date: 03 August 2000 (03.08.2000)
(21) International Application Number: PCT/NZ99/00214 (22) International Filing Date: 10 December 1999 (10.12.1999) (30) Priority Data: 333992 29 January 1999 (29.01.1999) NZ (60) Parent Application or Grant NEW ZEALAND INSTITUTE FOR CROP & FOOD RESEARCH LIMITED [/]; (). LISTER, Carolyn, Elizabeth [/]; (). EADY, Colin, Charles [/]; (). LISTER, Carolyn, Elizabeth [/]; (). EADY, Colin, Charles [/]; (). HAWKINS, Michael, Howard ; ().		Published
(54) Title: TRANSFORMATION AND REGENERATION OF ALLIUM PLANTS (54) Titre: TRANSFORMATION ET REGENERATION DE PLANTES APPARTENANT A LA FAMILLE ALLIUM (57) Abstract <p>The invention relates to a novel transformation method for plants of the genus Allium, in particular to onion plants. Plants transformed by the method are also provided. The method preferably involves an Agrobacterium tumefaciens-mediated transformation, more preferably involving immature embryos as the explant source and employing a binary vector.</p> (57) Abrégé <p>La présente invention concerne un nouveau procédé permettant de transformer des plantes du genre Allium, plus particulièrement des oignons et des plantes transformées au moyen dudit procédé. Ce procédé implique de préférence une transformation fondée sur Agrobacterium tumefaciens et implique plus préférentiellement encore l'utilisation d'embryons immatures en tant que source d'explant et d'un vecteur binaire.</p>		

PCTWORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁷ : C12N 15/82, A01H 5/00, 5/06	A1	(11) International Publication Number: WO 00/44919 (43) International Publication Date: 3 August 2000 (03.08.00)
(21) International Application Number: PCT/NZ99/00214 (22) International Filing Date: 10 December 1999 (10.12.99) (30) Priority Data: 333992 29 January 1999 (29.01.99) NZ (71) Applicant (for all designated States except US): NEW ZEALAND INSTITUTE FOR CROP & FOOD RESEARCH LIMITED [NZ/NZ]; Gerald Street, Lincoln, Christchurch (NZ). (72) Inventors; and (75) Inventors/Applicants (for US only): LISTER, Carolyn, Elizabeth [NZ/NZ]; 58 Charles Upham Avenue, Hillmorton, Christchurch (NZ). EADY, Colin, Charles [NZ/NZ]; 69 Ellesmere Junction Road, Lincoln, Christchurch (NZ). (74) Agents: HAWKINS, Michael, Howard et al.; Baldwin Shelston Waters, NCR Building, 342 Lambton Quay, Wellington (NZ).	(81) Designated States: AE, AL, AM, AT, AT (Utility model), AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, CZ (Utility model), DE, DE (Utility model), DK, DK (Utility model), DM, EE, EE (Utility model), ES, FI, FI (Utility model), GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SK (Utility model), SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG). Published With international search report.	
(54) Title: TRANSFORMATION AND REGENERATION OF <i>ALLIUM</i> PLANTS (57) Abstract <p>The invention relates to a novel transformation method for plants of the genus <i>Allium</i>, in particular to onion plants. Plants transformed by the method are also provided. The method preferably involves an <i>Agrobacterium tumefaciens</i>-mediated transformation, more preferably involving immature embryos as the explant source and employing a binary vector.</p>		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakhstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

Description

5

10

15

20

25

30

35

40

45

50

55

5

Transformation and regeneration of *Allium* plants

10

Field of Invention

- 5 The invention relates to a method of transforming plants of the *Allium* family and more particularly to the transformation of onion plants. The invention also relates to the transformed plants.

15

Background of the Invention

20

There are no published protocols for the transformation and regeneration of *Allium* species. The *Allium* crop species are probably the most economically important vegetable species for which transformation technology is unavailable. For other major vegetable crops, confirmed transformation systems have been produced.

15

25

Initially, many monocotyledons were thought to be unsusceptible to *Agrobacterium*-mediated transformation. The development of direct gene transfer techniques soon led to bombardment being the favoured method of monocotyledon transformation. However, direct gene transfer is not without its problems. Often, low transformation frequencies and a high frequency of unusual integration patterns has been observed in transgenic plants. Recently, *Agrobacterium*-mediated transformation of monocotyledons has gained favour and many monocotyledonous species (including rice, wheat, barley, maize and sugarcane) have now been transformed using this method. A key component in the success of these systems has been the transfer of DNA to callus cell types (usually derived from the pre culture of embryo tissue) followed by regeneration from these callus cells using precise post transformation selection protocols. Transformation of *Allium* callus is not useful as regeneration from callus is extremely difficult.

30

20

35

25

40

30

Recently, Haseloff (1997) has modified the *gfp* gene to enhance its use as a transgenic marker gene in viable plant systems. Green fluorescent protein (GFP) enables researchers to follow precisely the fate of any cells expressing this gene and

45

50

55

so optimise post transformation cell survival. Such a system has been useful in the development of the onion transformation protocol reported here.

As monocotyledonis, the *Allium* species were predisposed to be recalcitrant to transformation. Onions (*Allium cepa* L) are a crop with diverse environmental requirements. It has, therefore, been relatively understudied with respect to the application of biotechnology. There are only a few reports of DNA delivery to *Alliums* (Klein 1987; Dommissie et al. 1990; Eady et al. 1996; Barandiaran et al. 1998). Three workers used direct gene transfer whilst Dommissie et al.(1990) demonstrated that *Agrobacterium*-mediated transformation may be possible. Recently some reports of regeneration protocols for *Alliums* that are appropriate for transformation study have become available (Hong and Deberg 1995; Xue et al. 1997; Eady et al. 1998; Saker 1998). Only one report exists on the development of potential selective agents for use in *Allium* transformation (Eady and Lister 1998a).

Object of the Invention

It is therefore an object of the invention to provide a method for producing transgenic *Allium* plants or to at least provide the public with a useful choice.

In this specification we report the first repeatable protocol for the production of transgenic *Allium* plants.

Summary of the Invention

The invention provides a method of transforming *Allium* plants.

Preferably, the invention provides an *Agrobacterium tumefaciens* - mediated transformation method for *Allium* plants.

In particular, the invention provides a method of transforming plants of the *Allium* genus comprising inoculating an embryo culture of an *Allium* species with an *Agrobacterium tumefaciens* strain containing a suitable vector or plasmid.

5 Preferably embryos are inoculated immediately following their isolation.

Preferably the transformed plants are onions (*Allium cepa* L).

10 5 Preferably immature embryos are used as the explant source.

Preferably the embryos are transformed using a binary vector and more preferably a binary vector carrying a selectable gene.

10 The embryos may preferably be transformed with a herbicide selective gene. Examples include the *bar* gene or *ppt* gene encoding resistance to phosphinothricin or genes encoding resistance to glyphosate. However other genes may be used.

20 The embryos could alternatively be transformed with an antibiotic selective gene. An example is the kanamycin or geneticin resistance gene, *nptII*.

25 In particular, the invention provides a method of transforming *Allium* using immature embryos as an explant source, including:

- 30 20 a) isolating immature embryos of the *Allium* plant to be transformed;
b) innoculating the immature embryos with an *Agrobacterium tumefaciens* strain containing a binary vector;
c) wounding embryos and infiltrating embryos with *Agrobacteria*;
35 d) transferring embryos to a selective medium;
e) culturing embryo pieces;
25 f) selecting putative transgenic cultures; and
g) regenerating phenotypically normal plants.

40 The invention also provides transformed *Allium* plants. Preferably the *Allium* plants are transformed using protocols in line with the method of the invention.

30 45 **Brief Description of the Drawings**

Embodiments of the invention are now described, by way of example only, with

50

55

reference to the drawings in which:

Figure 1 shows a) GFP expression in embryo tissue after 5 days of cocultivation (x50). b) GFP expression after 2 weeks (x50). c) GFP sector after 6 weeks culture (x25). d) Independent GFP positive tissue (x5). e) GFP positive onion shoot culture (x5). f) Two GFP negative (left) and two GFP positive (right) roots from independent plants (x10). g) Transgenic onion plant (x0.2).

Figure 2 shows Southern analysis of the *gfp* gene of primary transformants: Bluescript plasmid containing the *gfp* gene (uncut), 1 copy number control (lane 1), 10 copy number control (lane 2), blank (lane 3), non transgenic onion (lane 4), 7 transgenic onion plants (lanes 5-11), bluescript plasmid containing the *gfp* gene (uncut), 1 copy number control (lane 12), 10 copy number control (lane 13), blank (lane 14), non transgenic onion (lane 15), 6 transgenic onion plants (lanes 16-21).

Figure 3 shows a Southern blot transgenic antisense root alliinase plants probed with the *gfp* gene fragment to indicate the presence of the pBINmgfpERantiroot T-DNA sequence. Lane 1 lambda hindIII marker; lane 2 one copy equivalent control pBINmgfpERantiroot, lane 3 five copy control pBINmgfpERantiroot; lane 4 non transformed onion, lane 5 positive control onion transformed with pBINmgfpER; lane 6-10 transgenic plants transformed with pBINmgfpERantiroot (6&7 and 9&10 are separate clones); lane 11 one copy equivalent control pBINmgfpERantiroot, lane 12 five copy control pBINmgfpERantiroot.

Figure 4 shows a Western blot analysis of alliinase levels in the roots of transgenic and non transgenic onion roots. Lane 1 purified root alliinase control; Lane 2-5 alliinase levels from the roots of four transgenic plants transformed with the pBINmgfpERantiroot DNA; Lane 6 alliinase levels from the roots of a typical non transgenic control plant.

Figures 5 and 6 show a Southern blot analysis of HindIII digested DNA from Onion plants transformed with the modified pCambia 3301 T-DNA. Figure 5 probed with *gfp* probe. Figure 6 probed with *bar* gene probe. Lane 1 and 2, 1 and 5 copy number controls of plasmid pBINmgfpER respectively (containing *gfp* gene), Lane 3 non-transgenic onion DNA. Lane 4-6 clones of a transformant selected from

experiment 994, Lane 7-9 clones of a transformant from experiment 9911. Lane 10-12 control transgenic plants containing the *gfp* gene and not the *bar* gene. Lane 13-14, 5 and 10 copy controls of plasmid containing the *bar* gene (lane 11-14 over washed the *gfp* probed blot).

Figure 7 shows a comparison between A, an onion leaf containing the *bar* gene (two on left) and onion leaves without the *bar* gene (four on right) 10 days after painting with 0.5% v/v solution of the herbicide Buster and B, C and D showing an onion plant without the *bar* gene (left) and containing the *bar* gene (right) 0, 3 and 10 days respectively after spraying with 0.05% vv solution of the herbicide Buster.

Detailed Description of the Invention

Materials and Methods

Plant material: Field grown, open-pollinated umbels of *Allium cepa* L. were used as a source of immature embryos. Immature embryos were isolated as described by Eady et al. (1998).

Bacterial strain: *Agrobacterium tumefaciens* strain LBA4404 containing the binary vector *pBIN m-gfp-ER* (Haseloff 1997) or binary vectors derived from p Cambia series were used. Cultures were grown to log phase in LB media containing an appropriate antibiotic and then stored at -80°C in 1 ml aliquots containing 15% glycerol. Aliquots were used to inoculate 50 ml overnight cultures. The following morning cultures were replenished with an equal volume of LB containing antibiotic and 100 µM acetosyringone and grown for a further 4 hours. *Agrobacteria* were isolated by 10 minute centrifugation at 4500 rpm and resuspended in an equal volume of P5 (Eady and Lister 1998a) containing 200 µM acetosyringone.

Transformation procedure: Isolated immature embryos were isolated in groups of 20-40, cut into approximately 1mm sections and then transferred into 0.8 ml of *Agrobacteria* and vortexed for 30 seconds. Following this treatment, embryos were placed under vacuum (~ 20 in. Hg) for 30 minutes before blotting dry on sterile Whatman #1 filter paper and then transfer to solid P5 medium (Eady and Lister 1998) (~ 40 embryos per plate). After 6 days cocultivation, embryo pieces were transferred

to P5 plus 10 mg/l geneticin and 200 mg/l timentin or 5 mg/l of Basta (active ingredient phosphothricin) and 200 mg/l of timentin depending on which binary vector was used. These embryo pieces were cultured in the dark under the same conditions as described for the production of secondary embryos (Eady et al. 1998). Cultures were transferred to fresh medium every 2 weeks. After 3-4 transfers, growing material was transferred to P5 plus 25 mg/l geneticin or 5 mg/l Basta depending on the binary vector used and grown for a further 8 weeks. During this time pieces of putative transgenic tissue that were obviously actively growing were transferred to regeneration medium (Eady et al. 1998). Shoot cultures were maintained for 12 weeks and developing shoots were transferred to 1/2MS media (Murashige and Skoog 1962) plus 20 mg/l geneticin or 5 mg/l of Basta as appropriate to induce rooting. Rooted plants were either transferred to 1/2MS plus 120 g/l sucrose to induce bulbing or transferred to soil in the glasshouse (12 h 12-23°C day, 12h 4-16°C night).

Analyses for transformation: For GFP expression, tissue was examined by observation under a fluorescence microscope (excitation 475 nm, emission 510 nm Haseloff et al. 1997). Larger tissues with high levels of expression were observed using hand held "shirt pocket" fluorescent lanterns (Zelco industries inc., 630 So. Columbus Ave, Mt Vernon NY 10551-4445). *NptI* expression was determined by the ability of regenerating plantlets to form roots on 1/2MS containing geneticin, *bar* expression was determined by the ability of regenerating plantlets to form roots on 1/2 MS containing Basta.

DNA isolation was performed using a nucleon phytopure plant DNA extraction kit (Amersham Lifescience, Buckinghamshire, England). Southern analysis followed the method of Timmerman et al. (1993) and used PCR-amplified probes to confirm the presence of the *gfp*, *nptI* and *bar* genes. Genomic DNA from the onions was digested with *HindIII*, which cuts once in the middle of the T-DNA.

Cytology: Chromosome counts were made from the root tips of 2 primary transformants and followed the procedure of Grant et al. (1984).

Example 1*Transformation and characterisation of primary transgenic tissue*

After three days of cocultivation, single cells expressing GFP could be observed in tissue transformed with T-DNA containing the mgfpER gene. Attempts to count cells expressing GFP after 5 days were abandoned as the variation within treatments and between embryo pieces was huge, with many embryo pieces showing no fluorescence and some exhibiting hundreds of fluorescing cells (Fig. 1a). In the latter case, distinguishing between multicellular 'stable' transformation events and multiple adjacent single-celled 'transient' events was not possible. Thus, large biases in any measurement of initial transfer could have occurred. As an alternative, treatments were given an initial transfer rating: *** being excellent initial transfer (~20-30% of tissue pieces with > 20 GFP positive cells per plate), ** represented average initial transfer (5-20% of tissue pieces with some GFP positive cells per plate, and * being poor initial transfer (< 5% of tissue pieces with GFP positive cells per plate) (Table 1). Contamination was a problem in many experiments. Often whole experiments (data not shown) had to be abandoned due to contamination, much of which probably arose from infected embryos.

Table 1. Summary of 5 transformation experiments. * - poor, ** - average, *** - excellent initial transfer, see text for details. Numbers in brackets represent the percentage of transformants from uncontaminated embryos (a - represents the stage and treatments which were transferred to the wrong selective media for 4 days).

Expt	N°. of embryos	% of embryos contaminated	Initial transfer	N°. of multicellular GFP tissue pieces		Independent plants	Positive Southern#
				4wk	8wk		
1	~400	100	*	-	-	-	-
2	~360	40	***	52(16)a	15(4.6)	2 (0.6)	1 of 1 tested
3	~440	0	***	72 (16)	44 (10)	12 (2.7)	8 of 8 tested

4	~520	60	**	a	11 (2.4)	3 (0.7)	2 of 2 tested
5	~200	100	*	-	-	-	-

After 2 weeks on selection medium, embryo pieces transformed with pBINmgfpER construct were screened for GFP expression and only pieces containing fluorescing cells were maintained (Fig. 1b). The vast majority of fluorescing cells died over the following four weeks. Some fluorescing cells divided into multicellular clusters of up to ~50 before their ability to fluoresce gradually faded. One interpretation of this was that the transformed cells were still reliant on surrounding non-transgenic cells, which died due to selection pressure and could no longer support the transgenic cells. The number of stable transgenic sectors arising from different plates within experiments varied from 0 to 21 and reflected the numerous parameters that affect the onion transformation process. Comparison between experiments was initially possible and ranged from * in contaminated samples to *** in non-contaminated samples. Indeed, lack of good initial transfer was often an early indication of contamination. Eady et al. (1998) and Eady and Lister (1998ab) demonstrated that genotype, condition of the embryo, size of the embryo, cocultivation conditions and selection pressure all affect embryo survival. The combined effects of these parameters and their interaction with the transformation process will, until they can be controlled, continue to make the success of onion transformation susceptible to large variation.

Example 2

Regeneration

After 6 weeks of culture, tissue was transferred to a selective medium without timentin. No growth of *Agrobacterium* was observed in any of the cultures grown on this medium. Fluorescing sectors continued to grow on this media and after 2 transfers it was possible to isolate the first sectors free from non-fluorescing cells (Fig. 1c). As sectors became independent (Fig. 1d) they were transferred to regeneration medium. A few sectors still attached to non-fluorescing tissue were also transferred. On regeneration medium transgenic cultures responded in the same way

as non-transgenic, embryo-derived cultures (Eady et al. 1998). Multiple shoots formed on many of the independent transgenic cultures. However, some, particularly the slower growing or more friable dedifferentiated cultures, either failed to regenerate or produced highly hyperhydric shoots that could not be transferred to the glasshouse.

Up to 29% of stable sectors produced shoot cultures from which plants could be obtained (Table 1). These responses to regeneration are typical of those seen in non-transformed embryogenic cultures (Eady et al. 1998). Actively growing shoots were transferred to rooting medium containing an appropriate selective agent. In the instances where non-fluorescing cells were also transferred to shoot media some shoots were produced that failed to root on geneticin. These did not fluoresce. All plants that formed actively growing roots on geneticin also fluoresced (Fig. 1f), indicating that in all instances the complete T-DNA was transferred. Fluorescence in the differentiated structures varied, with most fluorescence being seen in root tips. In shoots, strong fluorescence was limited to young shoots (Fig. 1e). However, GFP fluorescence in shoots was usually masked by red autofluorescence from the chlorophyll. The presence of GFP fluorescence in older leaves could sometimes be observed in the stomatal guard cells.

The multiple shoot cultures enabled clonal plants from independent transgenic events to be grown. This was particularly important as earlier attempts to exflask putative transgenic plants had failed (Eady and Lister 1998b). In the first successful transformation experiment described here only 4 from 48 transgenic plants transferred to the soil have died. A total of 14 independent transformants have been transferred to the containment glasshouse.

Example 3

Analyses of transformants from plants transformed with pBINmgfpER

Apart from fluorescence and growth on geneticin, transformation of onion plants was confirmed by Southern analysis, probing with the *gfp* gene (Fig. 2). As *HindIII* cuts the T-DNA only once it was possible to show copy number from the Southern analysis. Ten of the 13 transformants shown have single copies. The other 3 have 2 (lane 8), 3 (lane 18) and multiple copies (lane 7). Lanes 19 and 21 are from clonal shoots and,

as expected, they show the same pattern. *EcoR1* digest and subsequent Southern analysis liberated an expected internal T-DNA fragment of ~ 900 bp.

Chromosome counts in the 2 primary transformants tested showed a diploid ($2n = 16$) chromosome complement.

Example 4

Evidence that the transgenic onion plants transformed with the pBINmgfpERantiroot contain the antisense root alliinase gene construct.

Onion immature embryos were transformed according to the protocol of Eady et al (1999) with the *pBINmgfpER* plasmid (Haseloff 1997) modified to contain the antisense root alliinase gene construct. The BamH1 to Kpn1 fragment of the root alliinase gene was cloned into the BamH1 - Kpn1 sites in the cloning vector pART7. This gave a antisense version of the root alliinase sequence under control of the CaMV35s regulatory element and *ocs* termination sequence in pART7. The not1 fragment of this modified pART7 (containing the above *CaMV35s* promoter - antisense root alliinase - *ocs* termination) was then cloned into the Hind111 site of *pBINmgfpER*. Digestion of this plasmid (*pBINmgfpERantiroot*) with BamH1 to liberate a 1.6Kb fragment was used to determine presence and orientation of the insert. *pBINmgfpERantiroot* was electroporated into *Agrobacterium tumefaciens* strain LBA4404 and grown on kanamycin to select for transformants. Presence of the binary vector was confirmed by plasmid isolation and PCR for the *gfp* gene. LBA4404 (*pBINmgfpERantiroot*) was used in transformation experiments.

Six putative transformants that fluoresced (to indicate the presence of the *gfp* gene) and grew on media containing geneticin (to indicate presence of the *nptII* gene) were obtained from three experiments. Three of these transformants or clones thereof were analysed by Southern Blot analysis for the successful transfer of the T-DNA insert from the binary vector by probing with both *gfp* and *nptII* gene probes. Roots from these plants were also analysed biochemically for root alliinase enzyme activity following the protocol of Clark et al (1998) (Table 2). Western Blots of the desalted protein (0.5µg/lane) extracts were probed with an anti-alliinase antibody and visualised

colourmetrically using a goat-antirabbit-alkaline phosphatase system to determine the relative levels of alliinase enzyme in the transgenic plants.

Table 2:

Plant	Alliinase activity (U/mg protein)
Non transgenic CLK control (9910)	14.0
transformant 992.11F1	3.4
transformant 994.7G1	11.9
transformant 992.11F2	9.6
transformant 992.9A1	6.3

Results

1. Southern Analysis

All three plants analysed contained at least one copy of the T-DNA sequence containing the antisense root alliinase DNA sequence (Figure 3) indicating that integration of modified alliinase sequences into *Allium* species had been achieved. Both *nptII* and *gfp* sequences which flanked the antisense alliinase gene on the T-DNA could be deleted indicating successful transfer of the complete T-DNA in all cases.

The Western blot of Figure 4 shows the relative amounts of the root alliinase in protein extracts taken from the transgenic and control roots. These extracts were run on a 10% SDS page gel to separate the proteins and then transferred to nitrocellulose paper using standard techniques. This was then incubated with rabbit polyclonal antibodies raised against the purified alliinase (Clark S. A. 1993. Molecular cloning and cDNA encoding alliinase from onion (*Allium cepa* L.), Ph D. thesis, University of Canterbury, Christchurch, New Zealand). These antibodies have been shown to bind specifically to the alliinase protein. Goat anti-rabbit alkaline phosphatase was added to specifically bind this antibody and after washing, the membrane was immersed in NBT (4 nitroterazolium chloride) and BCIP (5 bromo 4 chloro 3 indolyl phosphate) for 30 minutes in the dark. Colour develops at the site of the phosphatase in proportion to the amount alliinase present. The Western blot therefore shows the relative amounts of alliinase protein present in the roots of the transgenic and control onion

plants. The control onion plant has the greatest colour development and has the most alliinase per unit of root protein. The intensity relates to the activity of the enzyme shown in the table and indicates that the activity is related to the amount of alliinase protein and not changes in enzyme activity. This is what is expected when using antisense technology to reduce enzyme activity.

Example 5

Transformation of onions to confer herbicide resistance.

Onion immature embryos were transformed according to the protocol with the plasmid pCambia3301 modified, using standard cloning techniques, to contain the *mgfpER* reporter gene construct instead of the *gus* reporter gene. This construct, contained within its T-DNA region the *bar* gene encoding resistance to the herbicide phosphinothricin and the visual reporter gene *mgfpER* both under regulatory control of the CaMV35s promoter. In two separate experiments, 994 and 9911 onion immature embryos from cultivar Cron 19 and CLK respectively were transformed with the above construct. In experiment 994 transformants were selected using the visual marker (GFP expression) and growth on herbicide, whilst on P5 media only. In experiment 9911 only selection on herbicide was used to select for transformation. Selection conditions in both experiments consisted of growth on P5 media containing 5mg/l of the herbicide Basta (containing phosphinothricin) and 200mg/l of timentin for 4-6 weeks with fortnightly subculture. Following this, cultures were transferred to P5 media plus 5mg/l Basta for a further 4 weeks of culture. Cultures were then transferred to SM4 media for 6 weeks. In experiment 9911 the SM4 media included 5mg/l Basta. Shoots from 9911 were rooted on 1/2MS30 plus 5mg/l Basta. Shoots from experiment 994 were just rooted in 1/2MS30. Once vigorous root growth was established plants were transferred to the glasshouse.

Results

Putative transgenic plants were produced from both experiments 3 from 994 and 4 from 9911. Southern Blot analysis of clones of one transformant from each experiment demonstrated that the *gfp* gene was present in plants from both experiments and that both cultivars could be transformed (Figure 5). When this blot was subsequently reprobbed for the presence of the *bar* gene (Figure 6) only the plants

selected solely on the basis of the herbicide resistance were shown to contain the *bar* gene. These plants were then used for herbicide leaf paint assays and subsequently sprayed with recommended field application rates of the herbicide Buster (active ingredient phosphinothricin). Control plants containing no *bar* gene showed noticeable wilt after 3 days and were essentially dead after 10 days following application of the herbicide, whilst the plants that contained the *bar* gene and had been selected on herbicide did not appear to be harmed and grew normally (Figure 7).

Example 6

Demonstration that transformation is independent of cultivar, construct, T-DNA and selective agent.

Further to examples 1-5, additional transgenic plants containing variations of the antisense alliinase construct (outlined in example 4) have been produced in additional cultivars CRON 12, CRON19 and CRON2. The nature of these plants has been confirmed by GFP expression where appropriate and regeneration on media containing geneticin. A summary of the plants produced in all the examples outlined above is given in table 3 to demonstrate the plasticity of the transformation system. In experiments that were not contaminated transgenic plants have been obtained from all cultivars so far tested.

Table 3 illustrating the different binary vectors, T-DNA, cultivars and selective agents used in the *Allium* transformation protocol outlined and the measures taken thus far to determine the nature of the transformants.

Experiment number	Binary vector used for transformation (T-DNA in brackets)	Cultivar used	Number of independent transformed lines produced	Number of plants produced	GFP +ve	Growth on selective agent (geneticin or Basta)	Southern Analysis and probe used
98/7,8,9	pBIN(mgfpER)	CLK	>10	>100	yes	yes (geneticin)	12/12 tested (gfp)
99/2	pBIN(antiroot-mgfpER)	CRON12	3	>40	yes	yes (geneticin)	5/5 tested (gfp & nptII)
99/3	pBIN(35santibulb-mgfpER)	CRON19	1	12	yes	yes (geneticin)	yet to be tested
99/3	pBIN(35santibulb-mgfpER)	CRON2	3	7	yes	yes (geneticin)	yet to be tested
99/4	pBIN(antiroot-mgfpER)	CRON19	1	3	yes	yes (geneticin)	yet to be tested
99/4	pCambia3301 (modified)	CRON19	1	10	yes	?(phosphinothricin)	3/3(gfp) tested 0/3(bar) tested
99/6	pCambia3301 (modified)	CRON12	1	2	yes	?(phosphinothricin)	Yet to be tested
99/6	pBIN(antiroot-mgfpER)	CRON12	1	1	yes	yes (geneticin)	yet to be tested
99/7	pBIN(mgfpER)	CRON19	1	>12	yes	yes (geneticin)	yet to be tested
99/11	pCambia3301 (modified)	CLK	3	>10	?	Yes (phosphinothricin)	3/3 (gfp & bar) tested
99/12	pBIN(bulbpromoterantibulb-mgfpER)	CLK	1	1	yes	yes	yet to be tested

(Basta = phosphinothricin)

Discussion

We have developed a repeatable transformation system for onion. The regenerating primary transformants appear to be phenotypically normal. The GFP expression, as a visual selectable marker, enabled post transformation selection conditions to be optimised. The GFP marker has also proved useful in the selection of transgenic plants from other species that are difficult to transform (Vain et al. 1998). Selection conditions have now been established, which enable the identification of transformants solely on their ability to root in selective media (example 5).

This method of producing transgenic onions is repeatable and efficient. It takes a short time to produce transgenic plants and utilizes techniques have been shown to be cultivar independent (Example 6).

For example, this described process of transformation can be used with any species within the *Allium* and is not limited to onions. Work has shown that the described process of transformation is genotype independent.

It is to be understood that the scope of the invention is not limited to the described embodiments and therefore that numerous variations and modifications may be made to these embodiments without departing from the scope of the invention as set out in this specification.

Industrial Applicability

The invention provides a novel method of transforming plants of the genus *Allium* and in particular onion plants. Also provided are plants transformed by the method. This allows *Allium* crop species which are an economically important vegetable species to be transformed by a variety of genes for improvement of *Allium* crop varieties.

References

- Barandiaran X, Dipietro A, Martin J (1998). *Plant Cell Reports* 17, 737-741.
- Dommissse EM, Leung DM, Shaw M L, Conner A J (1990). *Plant Science* 69, 249-257.
- Eady CC, Lister CE (1998a). *Plant Cell Reports* 18: 117-121.
- Eady CC, Lister CE (1998b). In: *Proceedings of the Second International Symposium on Edible Alliaceae*, Adelaide Australia. *Acta Horticulturae* In press
- Eady CC, Suo Y, Butler RC (1998). *Plant Cell Reports* 18: 111-116.
- Eady CC, Lister CE, Suo Y, Schaper D (1996). *Plant Cell Reports* 15, 958-962.
- Grant J, Brown A, Grace J (1984). *Australian Journal of Botany* 32, 665-677.
- Haseloff J, Siemering KR, Prasher DC, Hodge S (1997). *Proceedings of the National Academy of Sciences of the United States of America* 94, 2122-2127.
- Hong W, Debergh P (1995). *Plant Cell Tissue & Organ Culture* 43, 21-28.
- Klein TM, Wolf ED, Wu R, Sanford JC (1987). *Nature* 327, 70-73.
- Murashige T, Skoog F (1962). *Physiol. Plant* 15, 473-497.
- Saker MM (1998). *Biologia Plantarum* 40, 499-506.
- Timmerman GM, Fraw T J, Miller AL, Weeden NF, Jermyn WA (1993). *Theoretical*

- 5 *and Applied Genetics* **85**, 609-615.
- Vain P, Worland B, Kohli A, Snape JW, Christou P (1998). *Theoretical & Applied*
 Genetics **96**, 164-169.
- 10 Xue HM, Araki H, Kanazawa T, Harada T, Yakuwa T (1997). *Journal of the Japanese*
5 *Society for Horticultural Science* **66**, 353-358.

15

20

25

30

35

40

45

50

55

Claims

5

10

15

20

25

30

35

40

45

50

55

Claims

1. A method of transforming plants of the *Allium* genus.
2. A method according to claim 1 wherein the *Allium* genus is transformed with *Agrobacterium tumefaciens*.
3. A method of transforming plants of the *Allium* genus comprising inoculating an embryo culture of an *Allium* species with an *Agrobacterium tumefaciens* strain containing a suitable vector or plasmid.
4. A method according to any one of claims 1-3 in which the plants are onions.
5. A method according to any one of claims 1-4 wherein the embryos are transformed with a binary vector.
6. A method according to any one of claims 1-5 in which embryos of an *Allium* species are inoculated prior to their differentiation into callus tissue.
7. A method according to any one of claims 1-6 in which embryos of an *Allium* species are inoculated immediately following their isolation.
8. A method according to any one of claims 1-7 in which immature embryos are used.
9. A method of transforming *Allium* using immature embryos as an explant source, including:
 - (a) isolating immature embryos of the *Allium* plant to be transformed;
 - (b) innoculating cultures of the immature embryos with an *Agrobacterium tumefaciens* strain containing a binary vector;
 - (c) wounding embryos and infiltrating embryos with agrobacteria;
 - (d) transferring embryos to a selective medium;
 - (e) culturing embryo pieces;
 - (f) selecting putative transgenic cultures; and
 - (g) regenerating plants.
10. A method according to any one of claims 1-9 wherein the plant is transformed with an *Agrobacterium tumefaciens* strain containing a vector which carries a selectable gene.
11. A method according to claim 10 in which the selectable gene is a herbicide

- 5 resistance gene.
12. A method according to claim 11 in which the herbicide resistance gene is the *bar* gene or a glyphosate resistance gene.
- 10 13. A method according to claim 10 in which the selectable gene is an antibiotic resistance gene.
- 5 14. A method according to claim 13 in which the antibiotic resistance gene is the *nptII* gene.
- 15 15. A method according to any one of claims 1-14 wherein the plant is transformed with a modified alliinase gene.
- 10 16. A transformed plant produced by the method of any one of claims 1-15.
- 20 17. A transformed plant according to claim 16 which has an altered level of alliinase to that of an untransformed plant.




Figure 2




Figure 3




Figure 4




FIGURE 5




FIGURE 6





FIGURE 7



INTERNATIONAL SEARCH REPORT

International application No.

PCT/NZ99/00214

A. CLASSIFICATION OF SUBJECT MATTER												
Int. Cl. ⁷ : C12N 15/82, A01H 5/00, 5/06												
According to International Patent Classification (IPC) or to both national classification and IPC												
B. FIELDS SEARCHED												
Minimum documentation searched (classification system followed by classification symbols) DERWENT, CHEMICAL ABSTRACTS												
Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched BIOTECHDS, AGRICOLA												
Electronic data base consulted during the international search (name of data base and, where practicable, search terms used) Wpat, Chemical Abstracts, BiotechDS, Agricola: (Allium or onion or garlic) and transform? and agrobacter?												
C. DOCUMENTS CONSIDERED TO BE RELEVANT												
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.										
P,X	WO 99/10512 (NUNHEMS ZADEN) 4 March 1999 Page 13 line 24 to page 15 line 2, example 6	1-17										
X	Derwent Abstract Accession No. 92-060496/08, Class C06, D16 JP 04 004-879 (WAKUNAGA SEIYAKU KK) 9 January 1992 Whole Abstract	1										
X	WO 98/44136 (ZENCO (NO.4) LIMITED) 8 October 1998 Example 5 and claim 23	1-17										
<input checked="" type="checkbox"/> Further documents are listed in the continuation of Box C <input checked="" type="checkbox"/> See patent family annex												
* Special categories of cited documents: <table border="0"> <tr> <td>"A" document defining the general state of the art which is not considered to be of particular relevance</td> <td>"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention</td> </tr> <tr> <td>"E" earlier application or patent but published on or after the international filing date</td> <td>"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone</td> </tr> <tr> <td>"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)</td> <td>"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art</td> </tr> <tr> <td>"O" document referring to an oral disclosure, use, exhibition or other means</td> <td>"&" document member of the same patent family</td> </tr> <tr> <td>"P" document published prior to the international filing date but later than the priority date claimed</td> <td></td> </tr> </table>			"A" document defining the general state of the art which is not considered to be of particular relevance	"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention	"E" earlier application or patent but published on or after the international filing date	"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone	"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art	"O" document referring to an oral disclosure, use, exhibition or other means	"&" document member of the same patent family	"P" document published prior to the international filing date but later than the priority date claimed	
"A" document defining the general state of the art which is not considered to be of particular relevance	"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention											
"E" earlier application or patent but published on or after the international filing date	"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone											
"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art											
"O" document referring to an oral disclosure, use, exhibition or other means	"&" document member of the same patent family											
"P" document published prior to the international filing date but later than the priority date claimed												
Date of the actual completion of the international search 17 March 2000		Date of mailing of the international search report 22 MAR 2000										
Name and mailing address of the ISA/AU AUSTRALIAN PATENT OFFICE PO BOX 200, WODEN ACT 2606, AUSTRALIA E-mail address: pct@ipaustralia.gov.au Facsimile No. (02) 6285 3929		Authorized officer ANDREW ACHILLEOS Telephone No : (02) 6283 2280										

INTERNATIONAL SEARCH REPORT

International application No.

PCT/NZ99/00214

C (Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT		
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 92/06205 (CLOVIS MATTON N.V.) 16 April 1992 Page 2 lines 1-11, page 3 line 28 to page 4 line 35 and claim 2	1-17
X	WO 97/42333 (PIONEER HI-BRED INTERNATIONAL INC) 13 November 1997 Page 8 lines 15-20, page 13 lines 19-37	1-17

INTERNATIONAL SEARCH REPORT
Information on patent family members

International application No.
PCT/NZ99/00214

This Annex lists the known "A" publication level patent family members relating to the patent documents cited in the above-mentioned international search report. The Australian Patent Office is in no way liable for these particulars which are merely given for the purpose of information.

Patent Document Cited in Search Report				Patent Family Member			
WO	99/10512	AU	92647/98				
WO	98/44136	AU	74285/98	GB	9706469		
WO	97/42333	AU	29377/97				
WO	92/06205	CA	2089072	EP	554273	NL	9002116
END OF ANNEX							